

arXiv:math/9509226v1 [math.LO] 15 Sep 1995

WAS SIERPINSKI RIGHT III?
CAN CONTINUUM-C.C. TIMES
C.C.C. BE CONTINUUM-C.C.?
SH481

SAHARON SHELAH

Institute of Mathematics
The Hebrew University
Jerusalem, Israel

Rutgers University
Department of Mathematics
New Brunswick, NJ USA

ABSTRACT. We prove the consistency of: if B_1, B_2 are Boolean algebras satisfying the c.c.c. and the $2^{\aleph_0}$ -c.c. respectively then $B_1 \times B_2$ satisfies the $2^{\aleph_0}$ -c.c.

We start with a universe with a Ramsey cardinal (less suffice).

Typed 5/92 - Latest Revision 1/13/95

I thank Alice Leonhardt for the beautiful typing

§1 corrected 4/94

Partially supported by the basic research fund, Israeli Academy

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

§0 INTRODUCTION

We heard the problem from Velickovic who got it from Todorcevic, it says “are there P , a c.c.c. forcing notion, and Q is a $2^{\aleph_0}$ -c.c. forcing such that $P \times Q$ is not $2^{\aleph_0}$ -c.c.?” We can phrase it as a problem of cellularity of Boolean algebras or topological spaces.

We give a negative answer even for $2^{\aleph_0}$ regular, this by proving the consistency of the negation. The proof is close to [Sh 288], §3 which continues [Sh 276], §2 and is close to [Sh 289]. A recent use is [Sh 473].

We start with $V \models “\lambda$ is a Ramsey cardinal”, use c.c.c. forcing blowing the continuum to λ . Originally the paper contained the consistency of e.g. $2^{\aleph_0} \rightarrow [\aleph_2]_3^2, 2^{\aleph_0}$ the first k_2^2 -Mahlo, (weakly inaccessible)(remember $k_2^2 < \omega$) but the theorem presented arrive here to satisfactory state (for me) earlier. See more [Sh 546]. I thank Mariusz Rabus for corrections.

* * *

What problems do [Sh 276], [Sh 288], [Sh 289], [Sh 473] and [Sh 481] raise? The most important are (we state the simplest uncovered case for each point):

- A Question.** 1) Can we get e.g. $CON(2^{\aleph_0} \rightarrow [\aleph_2]_3^2)$; more generally raise μ^+ to higher.
 2) Can we get $CON(\aleph_\omega > 2^{\aleph_0} \rightarrow [\aleph_1]_3^2)$; generally lower 2^μ , the exact $\aleph_n$ seems to me less exciting.
 3) Can get e.g. $CON(2^\mu > \lambda \rightarrow [\mu^+]_3^2)$?

Also concerning [Sh 473].

- B Question.** 1) Can we get the continuity on a non-meagre set for functions $f : \kappa^2 \rightarrow \kappa^2$?
 2) what can we say on continuity of 2-place functions?
 3) What about n -place functions (after [Sh 288]).

- C Question.** 1) Can we get e.g. for $\mu = \mu^{<\mu} > \aleph_0, CON(\text{if } P \text{ is } 2^\mu\text{-c.c., } Q \text{ is } \mu^+\text{-c.c. then } P \times Q \text{ is } 2^\mu\text{-c.c.})$?
 2) Can we get e.g. $CON(\text{if } P \text{ is } 2^{\aleph_0}\text{-c.c., } Q \text{ is } \aleph_2\text{-c.c. then } P \times Q \text{ is } 2^{\aleph_0}\text{-c.c.})$
 3) Can we get e.g. $CON(2^{\aleph_0} > \lambda > \aleph_0, \text{ and if } P \text{ is } \lambda\text{-c.c., } Q \text{ is } \aleph_2\text{-c.c. then } P \times Q \text{ is } \lambda\text{-c.c.})$

On A1 see [Sh 546].

Discussion Maybe the solution to (A1) is by using squared demand and if in $\delta < \lambda, cf(\delta) > \mu$ we guess $\langle N_s : s \in [B]^{\leq 2} \rangle, \mathcal{C}$, try to by Q_δ to add a large subset of B on which only two colours appear; but we want to do it also when $otp(B) > \mu^+$. Naturally we assume that if $\delta' < \delta, cf(\delta') > \mu^+, \delta' = \sup(B \cap \delta')$ this was done $\langle N_s : s \in [B \cap \delta']^{\leq 2} \rangle$, but we need more: including dividing B to μ set on each only two colours (by Q_δ). To do this and have λ, k_3^2 -Mahlo (rather than measurable or $(\lambda \rightarrow^+ (\omega_1)_2^{<\omega})$) we have to use a very strong diamond.

For problem (A2) the natural thing is to use systems $\bar{N} = \langle N_s : s \in [B]^{\leq 2} \rangle$ which are not end extension systems. Then it is natural to use the forcing on this

stronger; “specializing” not only the colouring but all $P_{N_s \cap \lambda}$ -names of ordinals (as defined in §8). This required a suitable squared diamond; this has not yet been clarified (actually we need somewhat less than $\bar{N}, S$).

But for problem (A3) a weaker version of this suggest itself. As in the solution of 1, λ is k_2^2 -Mahlo $\langle \langle N_s^\delta : s \in [B^\delta]^{\leq 2} \rangle : \delta \in S \rangle$ is such that $N_s^\delta \subseteq H(\delta^+)$, (we think of N_s^δ as guessing the isomorphism type over $H(\delta)$). Now we have to define a preliminary forcing R , λ -complete or at least strategically λ -complete, satisfying the λ^+ -c.c. So we have “copies” of $\langle N_s^\delta : s \in [B^\delta]^{\aleph_2} \rangle$ which behave like Δ -systems. [Saharon].

But if want to get tree like systems (ease requirement on forcing) we need more (enough dependency). For simplicity $\lambda = cf(\chi) = \chi^\lambda$ and use the following instead forcing and do it with.

We can have in V (or force),

$\bar{C} = \langle C_\delta : \delta \in S \rangle$ a square, $S \subseteq \chi$, $\bar{S} =: \{\delta \in S : cf(\delta) = \lambda\} \subseteq \chi$ stationary, $[\delta \in S \Rightarrow otp(C_\delta) \leq \lambda]$, we have squared diamond $\bar{C} = \langle C_\delta : \delta \in S \rangle$, and we choose for $\delta \in S, B_\delta \prec \mathfrak{A}_\delta, \|B_\delta\| \leq |\omega + otp(C_\delta)|, \delta_1 \in C_{\delta_2} \Rightarrow B_{\delta_1} \prec B_{\delta_2}$ & $\langle B_\delta : \delta \in C_\delta \cup \{\delta_1\} \rangle \in B_{\delta_1}, \delta_2 = \sup C_{\delta_1} = B_{\delta_1} = \bigcup_{\delta \in C_{\delta_1}} B_\delta^*$.

Now we can copy the squared diamond $\langle \langle N_{\delta,s} : s \in [B_\delta]^{\leq 2} \rangle : \delta < \lambda \rangle$

getting $\langle \langle N_{\delta,s}^* : s \in [B_\delta^*]^{\leq 2} \rangle : \delta \in S \rangle$. We then define $\langle P_i, Q_i, A_i : i < \chi \rangle, |a_i| \leq \chi_2$ (or $|a_i| < \kappa$).

Concerning (B1) the expected theorem holds. For 2-place function, note that the Sierpinski colouring can be viewed as a function from ${}^\mu 2$ to $\{0_\mu, 1_\mu\} \subseteq {}^\mu 2$. So the best we can hope for is

(B2)' can we get the consistency of $(*)_\mu$ for any 2-place function f from ${}^\mu 2$ to ${}^\mu 2$ there are (everywhere) non-meagre $A \subseteq {}^\mu 2$ and continuous functions f_0, f_2 from A to ${}^\mu 2$ such that $(\forall \eta, \nu \in A)[f(\eta, \nu) \in \{f_0(\eta, \nu), f_1(\eta, \nu)\}]$.

So we have to put together the proofs of [Sh 473] (continuity on non-meagre), [Sh 288] ($2^{\aleph_0} \rightarrow [\aleph_1]_3^2$), using the k_2^2 -Mahlo only and replace $\aleph_0$ by μ .

For problem (B3), $\mu = \aleph_0$ we have to generalize [Sh 288], §3. But also for $\mu > \aleph_0$, we have to consider what can be said on the partition of trees (see [Sh 288], §4 for a positive answer for large cardinal (indestructible measurable $n^* < \omega$)).

Concerning (C2), the problem with the approach to (A1) is “why should Q_δ from [Sh 481], 1.7 satisfies Q_δ^2 is c.c.c.”

Similarly (C3)(A3). A natural approach is to consider $\langle N_s : s \in [B]^{< \aleph_0} \rangle$ and use a subset $X \in [B]^{otp B}$ such that for different uses we use almost disjoint X 's. This was not completed but we restrict ourselves to “not only P satisfies the $2^{\aleph_0}$ -c.c. but even P^n (for each $n < \omega$).

Concerning (C1) we cannot replace ${}^\mu n$ elements of B by 1, but we can use a directed system, so “ P satisfies the 2^μ -c.c.”, is replaced by “for $\sigma < \mu, P^\sigma$ satisfies the 2^μ -c.c.” (or slightly less).

Another question is Velickovic's question answered for Borel c.c.c. forcing in [Sh 480]; i.e. (C4).

Preliminaries

0.A. Let $<^*_\chi$ be a well ordering of

$H(\chi) = \{x : \text{the transitive closure of } x \text{ has cardinality } < \chi\}$ agreeing with the usual well ordering of the ordinals,

P (and Q, R) will denote forcing notions, i.e. partial order with a minimal element $\emptyset = \emptyset_P$.

A forcing notion P is λ -closed if every increasing sequence of members of P , of length less than λ , has an upper bound.

0.B. For sets of ordinals, A and B , define $H_{B,A}^{OP}$ as the maximal order preserving bijection between initial segments of A and B , i.e., it is the function with domain $\{\alpha \in A : otp(\alpha \cap A) < otp(B)\}$ and $H_{A,B}^{OP}(\alpha) = \beta$ if and only if $\alpha \in A$, $\beta \in B$ and $otp(\alpha \cap A) = otp(\beta \cap B)$.

Definition 0.1. $\lambda \rightarrow^+ (\alpha)_\mu^{<\omega}$ holds provided that: if whenever F is a function from $[\lambda]^{<\omega}$ to λ , $F(w) < \min(w)$, $C \subseteq \lambda$ is a club then there is $A \subseteq C$ of order type α such that $\left[w_1, w_2 \in [A]^{<\omega}, |w_1| = |w_2| \Rightarrow F(w_1) = F(w_2) \right]$.
(See [Sh:f], XVII, 4.x).

0.1A Remark. 1) If λ is Ramsey cardinal then $\lambda \rightarrow^+ (\lambda)_\mu^{<\omega}$.

2) If $\lambda = \text{Min}\{\lambda : \lambda \rightarrow (\alpha)_\mu^{<\omega}\}$ then λ is regular and $\lambda \rightarrow^+ (\alpha)_\mu^{<\omega}$.

Definition 0.2. $\lambda \rightarrow [\alpha]_{\kappa, \theta}^n$ if for every function F from $[\lambda]^n$ to κ there is $A \subseteq \lambda$ of order type α such that $\{F(w) : w \in [A]^n\}$ has power $\leq \theta$.

Definition 0.3. A forcing notion P satisfies the Knaster condition (has property K) if for any $\{p_i : i < \omega_1\} \subseteq P$ there is an uncountable $A \subseteq \omega_1$ such that the conditions p_i and p_j are compatible whenever $i, j \in A$.

§1 CONSISTENCY OF “C.C.C. $\times 2^{\aleph_0}$ -C.C. = $2^{\aleph_0}$ - C.C.”

The a_i 's are not really necessary but (hopefully) clarify.

1.1 Definition. 1) $\mathcal{K}_{\mu,\kappa}$ is the family of $\bar{Q} = \langle P_\gamma, \underline{Q}_\beta, a_\beta : \gamma \leq \alpha, \beta < \alpha \rangle$, where:

- (a) $\langle P_\gamma, \underline{Q}_\beta : \gamma \leq \alpha, \beta < \alpha \rangle$ is a finite support iteration
- (b) every $P_\gamma, \underline{Q}_\gamma$ satisfies the c.c.c.
- (c) $\underline{Q}_\beta$ is a P_β -name which depends just on $G_{P_\beta} \cap P_{a_\beta}^*$ (see below; hence it is in $V[G_{P_\beta}^*]$), and $|\underline{Q}_\beta| \leq \kappa$ and its set of members $\subseteq V$ (for simplicity)
- (d) $a_\beta \subseteq \beta$, $|a_\beta| \leq \mu$ and $\gamma \in a_\beta \Rightarrow a_\gamma \subseteq a_\beta$.

2) For such $\bar{Q}$ we call $a \subseteq \ell g(\bar{Q}), \bar{Q}$ -closed if $[\beta \in a \Rightarrow a_\beta \subseteq a]$ and let

$$P_a^* = P_a^{\bar{Q}} =: \left\{ p \in P_\alpha : \text{Dom}(p) \subseteq a \text{ and for all } \beta \in \text{Dom}(p) : p(\beta) \in V \right. \\ \left. (\text{not a name}) \text{ and } p \restriction a_\beta \Vdash “p(\beta) \in Q_\beta” \right\}$$

(so we are defining P_a^* by induction on $\sup(a)$) ordered by the order of $P_{\sup(a)}$.

3) $\mathcal{K}_{\mu,\kappa}^k$ is the class of $\bar{Q} \in \mathcal{K}_{\mu,\kappa}$ such that if $\beta < \gamma \leq \ell g(\bar{Q}), cf(\beta) \neq \aleph_1$ then P_γ/P_β satisfies the Knaster condition (actually we can use somewhat less). Let $\mathcal{K}_{\mu,\kappa}^n = \mathcal{K}_{\mu,\kappa}$.

4) If defining $\bar{Q}$ we omit P_α we mean $\bigcup_{\beta < \alpha} P_\beta$ if α is limit, $P_\beta * \underline{Q}_\beta$ if $\alpha = \beta + 1$.

5) We do not lose, if we assume $\underline{Q}_\beta \subseteq [\kappa]^{<\aleph_0}$ and the order $\subseteq$; (then 1.2(1)(g) becomes trivial as for closed $p, q \in P_j^*, p \restriction a \leq q \restriction a$).

1.2 Claim. 1) Assume $x \in \{n, k\}$ and $\bar{Q} = \langle P_\gamma, \underline{Q}_\beta, a_\beta : \beta < \alpha, \gamma \leq \alpha \rangle \in \mathcal{K}_{\mu,\kappa}^x$.

Then

- (a) for $\alpha^* < \alpha$, $\bar{Q} \restriction \alpha^* =: \langle P_\gamma, \underline{Q}_\beta, a_\beta : \beta < \alpha^*, \gamma \leq \alpha^* \rangle$ belongs to $\mathcal{K}_{\mu,\kappa}^x$
- (b) P_α^* is a dense subset of P_α
- (c) for any $\bar{Q}$ -closed $a \subseteq \alpha$, $P_a^* \leq P_\alpha$ (in particular P_α^* is a dense subset of P_α); moreover, if $p \in P_\alpha^*$ then $p \restriction a \in P_a^*$ and $[p \restriction a \leq q \in P_a^* \Rightarrow r =: q \cup p \restriction (\alpha \setminus a) \in P_\alpha^* \ \& \ p \leq r \ \& \ q \leq r]$
- (d) for a $\bar{Q}$ -closed $a \subseteq \alpha$, $\langle P_{a \cap \gamma}^*, \underline{Q}_\beta, a_\beta : \beta \in a, \gamma \in a \rangle$ belongs to $\mathcal{K}_{\mu,\lambda}^x$ (except renaming; not used)
- (e) if $\underline{Q}_\alpha$ is a P_a^* -name of a c.c.c. forcing notion of cardinality $\leq \kappa$, each member of $\underline{Q}_\alpha$ is from V , $a \subseteq \alpha$ is $\bar{Q}$ -closed, $|a| \leq \mu$ and $P_{\alpha+1} = P_\alpha * \underline{Q}_\alpha$ and $\underline{Q}_\alpha$ satisfies the Knaster condition or at least $\beta < \alpha \Rightarrow P_\alpha * \underline{Q}_\alpha / P_{\beta+1}$ satisfies the Knaster condition then $\langle P_\gamma, \underline{Q}_\beta, a_\beta : \beta < \alpha + 1, \gamma \leq \alpha + 1 \rangle \in \mathcal{K}_{\mu,\lambda}^x$

(f) if $n < \omega, p_1, \dots, p_n \in P_{\alpha^*}$ and

(*) for every $\beta \in \bigcup_{\ell=1}^n \text{Dom}(p_\ell)$ for some $m = m_{\beta, \ell} \in \{1, \dots, n\}$ we have
 $p_m \restriction \beta \Vdash "p_\ell(\beta) \leq_{Q_\beta} p_m(\beta) \text{ for } \ell = \{1, \dots, n\}"$

then $p_1, \dots, p_n$ has a least common upper bound p which is defined by:

$\text{Dom}(p) = \bigcup_{\ell=1}^n \text{Dom}(p_\ell), p_\ell(\beta) = p_{m_{\beta, \ell}}(\beta)$, so in particular $p \in P_{\alpha^*}$ and

$$\bigwedge_{\ell=1}^n p_\ell \in P_{\alpha^*} \Rightarrow p \in P_{\alpha^*}$$

(g) if $p_\ell \leq p$ and $p_\ell \in P_\gamma^*$ for $\ell < n$, and a_k is $\bar{Q}$ -closed for $k < m$ then there is $p' \in P_\gamma^*$, such that $p \leq p'$ and $P_{a_k}^* \models p_\ell \restriction a_k \leq p' \restriction a_k$ for $\ell < n, k < m$.

2) If $x \in \{n, k\}$ and $\delta < \lambda$ is a limit ordinal, for $\alpha < \delta$ we have

$\langle P_\gamma, Q_\beta, a_\beta : \beta < \alpha, \gamma \leq \alpha \rangle \in \mathcal{K}_{\mu, \lambda}^x$ and $P_\delta = \bigcup_{\gamma < \delta} P_\gamma$ then $\langle P_\gamma, Q_\beta, a_\beta : \beta < \delta, \gamma \leq \delta \rangle$

belongs to $\mathcal{K}_{\mu, \lambda}^x$.

Proof. Straightforward.

Essentially by [Sh 289], 2.4(2), p.176 (which is slightly weaker and its proof left to the reader, so we give details here).

1.3 Claim. Assume $\lambda \rightarrow^+ (\omega \alpha^*)_\mu^{<\omega}$ (e.g. λ a Ramsey cardinal, $\alpha^* = \lambda$) $\chi > \lambda$, $x \in H(\chi)$.

1) There is an end extension strong $(\chi, \lambda, \alpha^*, \mu, \aleph_0, \omega)$ -system for x (see Definition 1.3A).

2) There is an end extension $(\chi, \lambda, \alpha, \mu, \aleph_0, \omega)$ -system for x if x is Ramsey or $\lambda = \text{Min}\{\lambda : \lambda \rightarrow (\omega \alpha^*)_\mu^{<\omega}\}$ (also then the condition holds for every $\mu' < \mu$).

1.3A Definition. 1) We say $\bar{N} = \langle N_s : s \in [B]^{<1+n} \rangle$ is a $(\chi, \lambda, \alpha, \theta, \sigma, n)$ -system if:

- (a) $N_s \prec (H(\chi), \in)$ (or of some expansion)
 $\theta + 1 \subseteq N_s, \|N_s\| = \theta, {}^\sigma(N_s) \subseteq (N_s)$
- (b) $B \subseteq \lambda, \text{otp}(B) = \alpha$
- (c) $n \leq \omega$ (equally is allowed but $1 + \omega = \omega$ so s is always finite)
- (d) $N_s \cap N_t \subseteq N_{s \cap t}$
- (e) $N_s \cap B = s$
- (f) if $|s| = |t|$ then $N_s \cong N_t$ say $H_{s,t}$ is an isomorphism from N_s onto N_t (necessarily $H_{s,t}$ is unique)
- (g) if $s' \subseteq s, t' = \{\alpha \in t : (\exists \beta \in s') [|\beta \cap s| = |\alpha \cap t|]\}$ then $H_{s',t'}, H_{s,t}$ are compatible functions; $H_{s,s} = \text{id}$,
 $H_{s,t} \supseteq H_{s,t}^{OP}, H_{s_0, s_1} \circ H_{s_1, s_2} = H_{s_0, s_2}, H_{t,s} = (H_{s,t})^{-1}$
- (h) $\sup(N_s \cap \lambda) < \text{Min}\{\alpha \in B : \bigwedge_{\gamma \in s} \gamma < \alpha\}$.

2) We add the adjective “strong” if in strengthen clause (d) by

(d)⁺ $N_s \cap N_t = N_{s \cap t}$ (so in clause (g), $H_{s',t'} \subseteq H_{s,t}$).

3) We add the adjective “end extension” if

(i) $s \triangleleft t \Rightarrow N_s \cap \lambda \triangleleft N_t \cap \lambda$ (where $A \triangleleft B$ means $A = B \cap \min(B \setminus A)$)

4) We add “for x ” if $x \in N_s$ for every $s \in [B]^{<1+n}$, and $H_{s,t}(x) = x$.

1.3B Remark. If λ is a Ramsey cardinal (or much less see [Sh:f],XVII,4.x,[Sh 289],§4) then we have if $\gamma \in s \cap t$, $s \cap \gamma = t \cap \gamma$ and $y \in N_s$ then in $(H(\chi), \in, <_\chi^*)$ the elements y and $H_{t,s}(y)$ realizes the same type over $\{i : i < \gamma\}$. [prove?]

Proof. 1) Let $C = \{\delta < \lambda : \text{for every } \alpha < \delta \text{ there is } N \prec (H(\chi), \in, <_\chi^*) \text{ such that } \mu + 1 + \alpha \subseteq N \text{ and } \sup(N \cap \lambda) < \delta\}$. Clearly C is a club of λ .

Let $B_0 = \{\alpha_i : i < \omega\alpha^*\} \subseteq C$, $(\alpha_i \text{ strictly increasing})$ be indiscernible in $(H(\chi), \in, <_\chi^*, x)$ (see Definition 0.1). Let $B = \{\alpha_i : i < \omega\alpha^* \text{ limit}\}$. For $s \in [B_0]^{<\aleph_0}$ let N_s^0 = the Skolem Hull of $s \cup \{i : i \leq \mu\} \cup \{x, \lambda\}$ under the definable functions of $(H(\chi), \in, <_\chi^*)$ and

$$N_s = \bigcup \left\{ N_{t_1}^0 \cap N_{t_2}^0 : t_1, t_2 \in [\{\alpha_i : i < \omega\alpha^*\}]^{<\aleph_0} \text{ and } s = t_1 \cap t_2 \right\}.$$

Clearly

$$(*) \quad \|N_s\| \leq \mu \text{ and } \{x, \lambda\} \subseteq N_s.$$

Now we shall show

(*)₁ if $s \in [B]^{<\aleph_0}$, $y \in N_s$ then for every finite $t \subseteq B_0$ there is $u \in [B_0]^{<\aleph_0}$ such that $s \subseteq u$, $u^* \cap t \subseteq s$ and $y \in N_u^0$.

As $y \in N_s$ there are $s_1, s_2 \in [B_0]^{<\aleph_0}$ such that $y \in N_{s_1}^0 \cap N_{s_2}^0$ and $s = s_1 \cap s_2$. Let $s_1 \cup s_2 = \{\alpha_{i_0}, \dots, \alpha_{i_{m-1}}\}$ (increasing), and let $n^* = \sup\{n : \text{for some } \beta, \beta + n \in t\} + 1$, and define for $\ell \leq m$ a function f_ℓ with domain $s_1 \cup s_2$, such that

$$f_\ell(\alpha_{i_k}) = \begin{cases} \alpha_{i_{k+1}} & \text{if } k \geq m - \ell \text{ and } i_k \notin s \\ \alpha_{i_k} & \text{otherwise} \end{cases}$$

Note that

- ⊗₁ for $\ell < m$, $f_\ell \upharpoonright s_1 = f_{\ell+1} \upharpoonright s_1$ or $f_\ell \upharpoonright s_2 = f_{\ell+1} \upharpoonright s_2$ (or both)
[why? as $i_\ell \in s_2 \setminus s_1 \setminus s$ or $i_\ell \in s_2 \setminus s_1 \setminus s$ or $i_\ell \in s = s_1 \cap s_2$].
- ⊗₂ f_ℓ is order preserving with domain $s_0 \cup s_1$, $f_\ell \upharpoonright s$ = the identity.

As $y \in N_{s_1}^0 \cap N_{s_2}^0$ there are terms τ_1, τ_2 such that

$$y = \tau_1(\dots, \alpha_{i_k}, \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, \alpha_{i_k}, \dots)_{\alpha_{i_k} \in s_2}.$$

Using the indiscernibility of B_0 we can prove by induction on $\ell \leq m$ that

$$\otimes_{3,\ell} y = \tau_1(\dots, f_\ell(\alpha_{i_k}, \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, f_\ell(\alpha_{i_k}, \dots)_{\alpha_{i_k} \in s_2}.$$

[Why? For $\ell = 0$ this is given by the choice of τ_1, τ_2 . For $\ell + 1$ note that by $\otimes_2$, $f_{\ell+1} \circ f_\ell^{-1}$ is an order preserving function from $\text{Rang}(f_\ell)$ onto $\text{Rang}(f_{\ell+1})$.

By $\otimes_{3,\ell}$ and “ B_0 is indiscernible” we know

$\tau_1(\dots, f_\ell(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, f_\ell(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_2}$. By the last two sentences and the indiscernibility of B_0

$$\tau_1(\dots, (f_{\ell+1} \circ f_\ell^{-1})(f_\ell(\alpha_{i_k})), \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, (f_{\ell+1} \circ f_\ell^{-1})(f_\ell(\alpha_{i_k})), \dots)_{\alpha_{i_k} \in s_2}.$$

But $(f_{\ell+1} \circ f_\ell^{-1})(f_\ell(\alpha_{i_k})) = f_{\ell+1}(\alpha_{i_k})$ so

$$\tau_1(\dots, f_{\ell+1}(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, f_{\ell+1}(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_2}.$$

But by $\otimes_1$ for some $e \in \{1, 2\}$ we have $f_\ell \upharpoonright s_e = f_{\ell+1} \upharpoonright s_e$, so

$\tau_e(\dots, f_{\ell+1}(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_e} = \tau_e(\dots, f_\ell(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_e}$ but the latter is equal to y (by the induction hypothesis), hence the former so by the last sentence

$$y = \tau_1(\dots, f_{\ell+1}(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_1} = \tau_2(\dots, f_{\ell+1}(\alpha_{i_k}), \dots)_{\alpha_{i_k} \in s_2}.$$

So we have carried the induction on $\ell \leq m$, and for $\ell = m$ we get $y \in N_{f_m(s_1)}^0$, but by the choice of n^* and f_m clearly $f_m(s_1) \cap t \subseteq s$, and we have proved $(*)_1$.

Now we can note

$(*)_2$ if $s \in [B]^{<\aleph_0}$ and $y_1, \dots, y_n \in N_s$ then for some $s_1, s_2 \in [B_0]^{<\aleph_0}$ we have:
 $s = s_1 \cap s_2$ and $y_1, \dots, y_n \in N_{s_1}^0 \cap N_{s_2}^0$.

[Why? We can find $u_1, \dots, u_n \in [B_0]^{<\aleph_0}$ such that $s \subseteq u_\ell, y_\ell \in N_{u_\ell}^0$ (as $y_\ell \in N_s$).

Now by $(*)_1$ for each $\ell = 1, 2, \dots, n$ we can find $v_\ell \in [B_0]^{<\aleph_0}$ such that $s \subseteq v_\ell, s = v_\ell \cap (\bigcup_{m=1}^n u_m)$ and $y_\ell \in N_{v_\ell}^0$. Let $u = \bigcup_{i=1}^n u_\ell, v = \bigcup_{\ell=1}^n v_\ell$, clearly $y_1, \dots, y_n \in N_u^0 \cap N_v^0$ and $u \cap v = s$, as required].

Now as we have Skolem functions $(*)_2$ implies

$(*)_3$ $N_s \prec (H(\chi), \in, <_\chi^*)$

Also trivially

$(*)_4$ $N_s^0 \prec N_s$ hence $\mu + 1 \subseteq N_s$

$(*)_5$ $s \subseteq t \Rightarrow N_s \prec N_t$.

Also

$(*)_6$ $N_{s_1} \cap N_{s_2} = N_{s_1 \cap s_2}$ for $s_1, s_2 \in [B]^{<\aleph_0}$.

[Why? The inclusions $N_{s_1 \cap s_2} \subseteq N_{s_1} \cap N_{s_2}$ follows from $(*)_5$; for the other direction let $y \in N_{s_1} \cap N_{s_2}$. By $(*)_1$ as $y \in N_{s_1}$ there is t_1 such that $s_1 \subseteq t_1 \in [B_0]^{<\aleph_0}, t_1 \cap (s_1 \cup s_2) = s_2$ and $y \in N_{t_1}^0$. By $(*)_1$, as $y \in N_{s_2}$ there is t_2 such that $s_2 \subseteq t_2 \in [B_0]^{<\aleph_0}, t_2 \cap (s_1 \cup s_2 \cup t_1) = s_1$ and $y \in N_{t_2}^0$. So $y \in N_{t_1}^0 \cap N_{t_2}^0$, but easily $t_1 \cap t_2 = s_1 \cap s_2$].

$(*)_7$ $\sup(N_s \cap \lambda) < \text{Min}\{\alpha \in B : \bigwedge_{\gamma \in s} \gamma < \alpha\}$.

[why? as $B_0 \subseteq C$ and see the Definition of C].

Now check that (a)-(h) of Definition 1.3A holds.

Now $\langle N_s : s \in [B]^{<\aleph_0} \rangle$ is as required.

2) If λ is Ramsey, without loss of generality $\text{otp}(B_0) = \lambda$ and it is easy to check 1.3A(i). The other case is like [Sh 289], §4. $\square_{1.3}$

1.4 Theorem. Assume $\aleph_0 < \mu \leq \kappa < \lambda = \text{cf}(\lambda)$, λ strongly inaccessible, λ a Ramsey cardinal, and $\diamond_{\{\delta < \lambda : \text{cf}(\delta) = \aleph_1\}}$ (can be added by a preliminary forcing). Then we have P such that:

- (a) P is a c.c.c. forcing of cardinality λ adding λ reals (so the cardinals and cardinal arithmetic in V^P should be clear), in particular in V^P we have $2^{\aleph_0} = \lambda$
- (b) $\Vdash_P$ “ MA holds for c.c.c. forcing notions of cardinality $\leq \mu$ and $< \lambda$ dense sets (and even for c.c.c. forcing notions of cardinality $\leq \kappa$ which are from $V[A]$ for some $A \subseteq \mu$)”
- (c) $\Vdash_P$ “if B is a λ -c.c. Boolean algebra, $x_i \in B \setminus \{0\}$ for $i < \lambda$ then for some $Z \subseteq \lambda$, $|Z| = \aleph_1$ and $\{x_i : i \in Z\}$ generates a proper filter of B (i.e. no finite intersection is 0_B)”
- (d) $\Vdash_P$ “if B_1 is a c.c.c. Boolean algebra, B_2 is a λ -c.c. Boolean algebra then $B_1 \times B_2$ is a λ -c.c. Boolean algebra.”

Proof. Let $\langle A_\delta : \delta < \lambda, \text{cf}(\delta) = \aleph_1 \rangle$ exemplifies the diamond. We choose by induction on $\alpha < \lambda$, $\bar{Q}^\alpha = \langle P_\gamma, \bar{Q}_\beta, a_\beta : \gamma \leq \alpha, \beta < \alpha \rangle \in \mathcal{K}_{\mu, \kappa}^n$ such that $\alpha^1 < \alpha \Rightarrow \bar{Q}^{\alpha^1} = \bar{Q}^\alpha \restriction \alpha^1$. In limits α use 1.2(2), for $\alpha = \beta + 1$, $\text{cf}(\beta) \neq \aleph_1$ take care of (b) by suitable bookkeeping using 1.2(1)(e). If $\alpha = \beta + 1$, $\text{cf}(\beta) = \aleph_1$ and A_β codes $p \in P_\beta$ and P_β -names of a Boolean algebra B_β and sequence $\langle x_i^\beta : i < \beta \rangle$ of non-zero members of B_β , and p forces ($\Vdash_{P_\beta}$) that there is in $V[G_{P_\beta}]$ some c.c.c. forcing notion Q of cardinality $\leq \mu$ adding some $Z \subseteq \beta$, $|Z| = \aleph_1$ with $\{x_i^\beta : i \in Z\}$ generating a proper filter of B_β then we choose $\bar{Q}_\beta$, if $p \in G_{P_\beta}$, as such Q . If $p \notin G_{P_\beta}$ or there is no such Q in $V[G_{P_\beta}]$, then $\bar{Q}_\beta$ is e.g. Cohen forcing.

So every $\bar{Q}^\alpha$ is defined, let $P = \bigcup_{\gamma < \lambda} P_\gamma$. Clearly (a) + (b) holds and (d) follows

by (c). So the rest of the proof is dedicated to proving (c).

So let $p \in P$, $p \Vdash$ “ $\bar{B}$ a λ -c.c. Boolean algebra, $x_i \in \bar{B} \setminus \{0_B\}$ for $i < \lambda$ ” without loss of generality the set of members of $\bar{B}$ is λ .

Let $x = \langle P, p, \bar{B}, \langle x_i : i < \lambda \rangle \rangle$, $\chi = \lambda^+$, by Claim 1.3 there are $A \in [\lambda]^\lambda$ and $\langle N_s : s \in [A]^{<\aleph_0} \rangle$ as there (for $\kappa = \mu + \kappa$ here standing for μ there). Let

$$C = \left\{ \delta < \lambda : \delta \text{ a strong limit cardinal } > \kappa + \mu, [\alpha < \delta \Rightarrow \bar{Q} \restriction \alpha \in H(\delta)], \right. \\ \delta = \sup(A \cap \delta), s \in [A \cap \delta]^{<\aleph_0} \Rightarrow \sup(\lambda \cap N_s) < \delta, \\ \left. \bar{B} \restriction \delta \text{ a } P_\delta\text{-name, and for } i < \delta \text{ we have } x_i \text{ a } P_\delta\text{-name} \right\}.$$

For some accumulation point δ of C , $cf(\delta) = \aleph_1$ and A_δ codes $\langle p, \underline{B} \restriction \delta, \langle \underline{x}_i : i < \delta \rangle \rangle$. We shall show that for some q , $p \leq q \in P_\delta$ and $q \Vdash_{P_\delta}$ “there is $\underline{Q}$ as required above”. By the inductive choice of $\underline{Q}_\delta$ this suffices.

Let $A^* \subseteq A \cap \delta$, $\text{otp}(A^*) = \omega_1$, $\delta = \sup(A^*)$ and $\langle \delta_i : i < \omega_1 \rangle$ increasing continuous, $\delta = \bigcup_{i < \omega_1} \delta_i$, $\delta_i \in C$, $A^* \cap \delta_0 = \emptyset$, $|A^* \cap [\delta_i, \delta_{i+1})| = 1$.

In V^{P_δ} we define:

$$\underline{Q} = \left\{ u : u \in [A^*]^{<\aleph_0}, \text{ and } \underline{B} \models \left(\bigcap_{i \in u} x_i \neq 0_{\underline{B}} \right) \right\}$$

ordered by inclusion. It suffices to prove that some q , $p \leq q \in P_\delta$, q forces that: $\underline{Q}$ is c.c.c. with $\bigcup \underline{G}_{\underline{Q}}$ an uncountable set; now clearly q forces that $\{x_i : i \in \bigcup \underline{G}_{\underline{Q}}\}$ generates a proper filter of $\underline{B}$.

If not, we can find q_i , u_i such that:

$$p \leq q_i \in P_\delta^* \text{ and } q_i \Vdash_{P_\delta} “u_i \in \underline{Q}” \text{ (where } u_i \in [A^*]^{<\aleph_0})$$

and $\langle (q_i, u_i) : i < \omega_1 \rangle$ are pairwise incompatible in $P_\delta * \underline{Q}$.

Let v_i be a finite subset of A^* such that: $u_i \subseteq v_i$, and

$$(*) \ [v \subseteq A^* \ \& \ v \text{ finite} \ \& \ \gamma \in (\text{Dom } q_i) \cap N_v \Rightarrow \gamma \in (\text{Dom } q_i) \cap N_{v \cap v_i}].$$

By Fodor’s Lemma for some stationary, $S \subseteq \omega_1$, u^* , v^* , n^* and $i(*)$ we have: for $i < j$ in S ,

$$v_i \cap \delta_i = v^* \subseteq \delta_{i(*)}, v_i \subseteq \delta_j, |v_i| = n^*, u_i \cap \delta_i = u^*$$

$$i(*) = \text{Min}(S)$$

$$\{|\gamma \cap v_i| : \gamma \in u_i\} \text{ does not depend on } i$$

$$q_i \restriction \delta_i \in P_{\delta_{i(*)}}^*$$

$$q_i \in P_{\delta_j}^*.$$

Let $b_i = N_{v_i} \cap \lambda$, so b_i is necessarily $\bar{Q}^\delta$ -closed and $|b_i| = \kappa$. Let $q_i^1 = q_i \restriction b_i$, so necessarily $q_i^1 \in P_{b_i}^*$ (see 2.2(1)(c)). Easily $P_{b_i}^* \subseteq N_{v_i}$ (though do not belong to it) so $q_i^1 \in N_{v_i}$.

Let $q_i^2 =: H_{v_{i(*)}, v_i}(q_i^1)$, so $q_i^2 \in P_{b_{i(*)}}^*$; let $q_i^3 =: (q_i \restriction \delta_{i(*)}) \cup \left[q_i^2 \restriction (N_{v_{i(*)}} \cap \lambda \setminus \delta_{i(*)}) \right]$ by 1.2(1)(c) we know $q_i^3 \in P_{\sup(b_{i(*)})+1}^*$ and $q_i^2 \leq q_i^3$, even without loss of generality $q_i^2 \leq q_i^3 \restriction b_{i(*)}$. As $P_{\sup b_{i(*)}+1}^* \triangleleft P_\delta$ and P_δ satisfies the c.c.c. clearly for

some $i < j$ from S , q_i^3, q_j^3 , are compatible in $P_{\sup(b_{i(*)})+1}^*$, so let $r \in P_{\sup(b_{i(*)})+1}^*$ be a common upper bound. So $q_i^3 \upharpoonright (\delta_{i(*)} \cap b_{i(*)}) \leq r \upharpoonright (\delta_{i(*)} \cap b_{i(*)})$ and $q_j^3 \upharpoonright (\delta_{i(*)} \cap b_{i(*)}) \leq r \upharpoonright (\delta_{i(*)} \cap b_{i(*)})$ and $q_i^3 \upharpoonright b_{i(*)} \leq r \upharpoonright b_{i(*)}, q_j^3 \upharpoonright b_{i(*)}$.

Without loss of generality $\text{Dom}(r) \subseteq b_{i(*)} \cup \delta_{i(*)}$ (allowed as $b_{i(*)}$ and $\delta_{i(*)}$ are closed, see 1.2(1)(c)); let $r_i = H_{v_i, v_{i(*)}}(r \upharpoonright b_{i(*)})$ and similarly $r_j = H_{v_j, v_{i(*)}}(r \upharpoonright b_{i(*)})$.

Note that $r_i \in P_{\delta_j}^*, r_j \in P_{\delta_i}^*, r_j \upharpoonright \delta_j = r_i \upharpoonright \delta_i = r \upharpoonright \delta_{i(*)}$. Hence $r_i \cup r_j \in P_{\delta}^*$.

Case 1. $r_i \cup r_j$ do not force (i.e. $\Vdash_{P_{\delta}}$) that

$$\bar{B} \models \text{“} \bigcap_{\alpha \in u_i \cup u_j} x_{\alpha} = 0_B \text{”}.$$

Then there is $r' \in P_{\delta}$, $r_i \leq r', r_j \leq r'$ forcing the negation. So without loss of generality $r' \in P_{\delta}^*$, and (as all parameters appearing in the requirements on r' are in $N_{v_i \cup v_j}$ also) $r' \in P_{\lambda \cap (N_{v_i \cup v_j})}^*$. Now

r', r, q_i, q_j has an upper bound $r'' \in P_{\delta}$.

[Why? By 1.2(1)(f), we have to check the condition (*) there, so let $\beta \in \text{Dom}(r') \cup \text{Dom}(r) \cup \text{Dom}(q_i) \cup \text{Dom}(q_j)$].

Subcase a. $\beta \in \delta_{i(*)} \setminus N_{v_i \cup v_j}$. Note that $N_{v_i \cup v_j} \cap \delta_{i(*)} = N_{v^*} \cap \lambda = b_{i(*)}$ (see choice of the N_u 's and definition of the b_{ε} 's) but $\text{Dom}(r') \subseteq N_{v_i \cup v_j} \cap \lambda$, so $\beta \notin \text{Dom}(r')$. Now

$$q_i \upharpoonright \delta_i = q_i \upharpoonright \delta_{i(*)} = q_i^3 \upharpoonright \delta_{i(*)} \leq r$$

$$q_j \upharpoonright \delta_j = q_j \upharpoonright \delta_{i(*)} = q_j^3 \upharpoonright \delta_{i(*)} \leq r.$$

So $r \upharpoonright \beta \Vdash_{P_{\beta}} \text{“} q_i(\beta) \leq r(\beta), q_j(\beta) \leq r(\beta) \text{”}$ and $\beta \notin \text{Dom}(r')$. So we have confirmed (*) from 1.2(1)(f) for this subcase.

Subcase b. $\beta \in \delta_{i(*)} \cap N_{v_i \cup v_j}$.

Exactly as above:

$N_{v_i \cup v_j} \cap \delta_{i(*)} = N_{v^*} \cap \lambda = b_{i(*)}$, so $\beta \in N_{v^*}, \beta \in \delta_{i(*)} \cap b_{i(*)}$. Also

$q_i \upharpoonright b_{i(*)} = q_i^1 \upharpoonright \delta_{i(*)} = q_i^2 \upharpoonright \delta_{i(*)} = q_i^3 \upharpoonright (\delta_{i(*)} \cap b_{i(*)}) \leq r \upharpoonright (\delta_{i(*)} \cap b_{i(*)})$ and

$q_j \upharpoonright b_{i(*)} = q_j^1 \upharpoonright \delta_{i(*)} = q_j^2 \upharpoonright \delta_{i(*)} = q_j^3 \upharpoonright (\delta_{i(*)} \cap b_{i(*)}) \leq r \upharpoonright (\delta_{i(*)} \cap b_{i(*)})$ and $r \upharpoonright (\delta_{i(*)} \cap b_{i(*)}) \leq r'$ (as $H_{v_i, v_{i(*)}}$ is the identity on $\delta_{i(*)} \cap b_{i(*)}$).

The last three inequalities confirm the requirement in 1.2(1)(f) (as $\beta \in \delta_{i(*)} \cap b_{i(*)}$, see above).

Subcase c. $\beta \in (\delta \setminus \delta_{i(*)}) \setminus N_{v_i \cup v_j}$.

In this case $\beta \notin \text{Dom}(r')$ (as $r' \in N_{v_i \cup v_j}$). Also $\delta_{i(*)} < \delta_i < \delta_j < \delta$ and:

$$\text{Dom}(r) \setminus \delta_{i(*)} \subseteq (b_{i(*)} \cup \delta_{i(*)}) \setminus \delta_{i(*)} \subseteq [\delta_{i(*)}, \delta_i)$$

$$\text{Dom}(q_i) \setminus \delta_{i(*)} \subseteq [\delta_i, \delta_j)$$

$$\text{Dom}(q_j) \setminus \delta_{i(*)} \subseteq [\delta_j, \delta).$$

So β belongs to at most one of $\text{Dom}(r'), \text{Dom}(r), \text{Dom}(q_i), \text{Dom}(q_j)$ so the requirement $(*)$ from 1.2(1)(f) holds trivially.

Subcase d. $\beta \in (\delta \setminus \delta_{i(*)}) \cap N_{v_i \cup v_j}$.

Clearly $\beta \notin \text{Dom}(r)$.

We know $q_i \restriction b_i = q_i^1, r_i \leq r', H_{v_i(*), v_i}(q_i^1) = q_i^2 \leq q_i^3 \restriction b_{i(*)} \leq r \restriction b_{i(*)}$ hence $q_i^1 \leq H_{v_i(*), v_i}^{-1}(r \restriction b_{i(*)}) = H_{v_i, v_i(*)}(r \restriction b_{i(*)}) = r_i$ but $r_i \leq r'$, so together $q_i^1 \leq r'$, and similarly $q_j^1 \leq r'$. As we have noted $\beta \notin \text{Dom}(r)$ we have finished confirming condition $(*)$ from 1.2(1)(f).

So really r, r', q_i, q_j has a least common upper bound, hence $(r'', u_i \cup u_j) \in P_\delta * Q$ exemplified $(q_i, u_i), (q_j, u_j)$ are compatible, as required.

Case 2. Not 1.

Let $\langle s_\beta : \beta < \lambda \rangle$ be such that:

$$s_\beta \in [A]^{<\aleph_0}, v^* \subseteq s_\beta, |s_\beta \setminus v^*| = |v_i \setminus v^*|, \sup(v^*) < \delta_{i(*)} < \min(s_\beta \setminus v^*)$$

$$\delta < \min(s_\beta \setminus v^*) \text{ (for simplicity)}$$

$$\beta < \gamma \Rightarrow \max(s_\beta) < \min(s_\gamma \setminus v^*).$$

As the truth value of $\bigcap_{\alpha \in u_i} x_\alpha$ is a P_a^* -name for some closed $a \in N_{v_i}$ of cardinality $\leq \mu$, and $q_i \restriction B \restriction \bar{B} \models \text{“} \bigcap_{\alpha \in u_i} x_\alpha \neq 0_{\bar{B}} \text{”}$ clearly $q_i^1 \restriction B \restriction \bar{B} \models \text{“} \bigcap_{\alpha \in u_i} x_\alpha \neq 0_{\bar{B}} \text{”}$.

For $\beta < \lambda$ let $\gamma^\beta = H_{s_\beta, v_i(*)}(r \restriction b_{i(*)})$, and $u'_\beta = H_{s_\beta, v_i(*)}(u_0)$. Let

$$Y = \{\beta < \lambda : r^\beta \in G_P\}.$$

Clearly:

$$r^\beta \restriction P_\lambda \restriction \bar{B} \restriction \bar{B} \models \text{“} \bigcap_{i \in u'_\beta} x_i \neq 0_{\bar{B}} \text{”}.$$

Clearly $p \leq r^\beta$ and for some β we have $r^\beta \restriction \bar{Y} \in [\lambda]^\lambda$ (and $p \in G_P$) and by the assumption of the case:

$$p \restriction \bar{B} \models \text{“} \left\{ \bigcap_{i \in u'_\beta} x_i : \beta \in Y \right\} \text{ is a set of non-zero members of } \bar{B} \text{ any two having zero intersection in } \bar{B} \text{”}.$$

This contradicts an assumption on B . $\square_{1.4}$

* * *

We can phrase the consistency result as one on colouring.

1.5 Lemma. 1) In 1.4 we can add:

- (e) if c is a symmetric function from $\left[2^{\aleph_0}\right]^{<\omega}$ to $\{0,1\}$ then at least one of the following holds:
- (α) we can find pairwise disjoint $w_i \subseteq 2^{\aleph_0}$ for $i < 2^{\aleph_0}$ such that:
 $c \upharpoonright [w_i]^{<\omega}$ is constantly zero but

$$\bigwedge_{i < j} (\exists u \subseteq w_i, \exists v \subseteq w_j) \left[c[u \cup v] = 1 \right]$$
 - (β) we can find an unbounded $B \subseteq 2^{\aleph_0}$ such that $c \upharpoonright [B]^{<\omega}$ is constantly 0.

It is natural to ask:

1.6 Question. Can we replace $2^{\aleph_0}$ by $\lambda < 2^{\aleph_0}$? $\aleph_1$ by $\mu < \lambda$? What is the consistency strength of the statements we prove consistent? (see later). Does λ strongly inaccessible k_2^2 -Mahlo (see [Sh289]) suffice?

1.7 Discussion. Of course, $1.5(e) \Rightarrow 1.4(c) \Rightarrow 1.4(d)$. Starting with λ weakly compact we can get a c.c.c. forcing notion P of cardinality λ , such that in V^P , $2^{\aleph_0} = \lambda$ and (e) of 1.5 holds for $c : \left[2^{\aleph_0}\right]^2 \rightarrow \{0,1\}$ (so $c(u) = 0$ if $|u| \neq 2$) and this suffices for the result. Also we can generalize to higher cardinals. We shall deal with this elsewhere.

1.8 Theorem. Concerning the consistency strength, in 1.4 it suffices to assume

- (*) λ is strongly inaccessible and for every $F : [\lambda]^{<\aleph_0} \rightarrow \mu$ and club C we can find $B \subseteq C$, (or just $B \subseteq \lambda$) $\text{otp}(B) = \omega_1$ such that
 - (a) B is F -indiscernible i.e. if $n < \omega, u, v \in [B]^n$ then $F(u) = F(v)$
 - (b) for every $n < \omega$ there is $B' \in [C]^\lambda$ such that:

$$\text{if } u \in [B']^n \text{ and } v \in [B]^n \text{ then } F(u) = F(v)$$

Proof. Let $R = \{\bar{Q} : \bar{Q} \in H(\lambda), \bar{Q} \in \mathcal{K}_{\mu,\kappa}^n\}$ ordered by $\bar{Q}^1 < \bar{Q}^2$ if $\bar{Q}^1 = \bar{Q}^2 \upharpoonright \ell g(\bar{Q}^1)$. Clause (b) takes care also of “the end extension” clause and for 1.3(A)(4), Clause (b) the proof is the same.

A somewhat less natural property though suffices.

(Note: Clause (b) also helps to get rid of the club C).

1.9 Claim. *In 1.4 it suffices to assume*

- (*)' *if $F : [\lambda]^{<\aleph_0} \rightarrow \mu$ then there is $B \subseteq \lambda$, $otp(B) = \omega$ such that*
- (a) *$F \upharpoonright [B]^n$ is constant for $n < \omega$*
 - (b) *if $u \triangleleft v^\ell \in [B]^{<\aleph_0}$ for $\ell = 1, 2$ then we can find $v_i \in [\lambda]^n$ for $i < \lambda$, $u \subseteq v_i$, $\min(v_i \setminus u) \geq i$, and $i < j \Rightarrow F(v^1 \cup v^2) = F(v_i \cup v_j)$.*

REFERENCES

- [Sh 546] Saharon Shelah. Was Sierpiński right? IV. *Journal of Symbolic Logic*, **accepted**.
- [Sh 276] Saharon Shelah. Was Sierpiński right? I. *Israel Journal of Mathematics*, **62**:355–380, 1988.
- [Sh 289] Saharon Shelah. Consistency of positive partition theorems for graphs and models. In *Set theory and its applications (Toronto, ON, 1987)*, volume 1401 of *Lecture Notes in Mathematics*, pages 167–193. Springer, Berlin-New York, 1989. ed. Steprans, J. and Watson, S.
- [Sh 288] Saharon Shelah. Strong Partition Relations Below the Power Set: Consistency, Was Sierpiński Right, II? In *Proceedings of the Conference on Set Theory and its Applications in honor of A.Hajnal and V.T.Sos, Budapest, 1/91*, volume 60 of *Colloquia Mathematica Societatis Janos Bolyai. Sets, Graphs, and Numbers*, pages 637–638. 1991.
- [Sh 480] Saharon Shelah. How special are Cohen and random forcings i.e. Boolean algebras of the family of subsets of reals modulo meagre or null. *Israel Journal of Mathematics*, **88**:159–174, 1994.
- [Sh 473] Saharon Shelah. Possibly every real function is continuous on a non-meagre set. *Publications de L’Institute Mathématique - Beograd, Nouvelle Série*, **57(71)**:47–60, 1995.
- [Sh 481] Saharon Shelah. Was Sierpiński right? III Can continuum–c.c. times c.c.c. be continuum–c.c.? *Annals of Pure and Applied Logic*, **78**:259–269, 1996.
- [Sh:f] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer, 1998.

PRIVATE APPENDIX: OLD §2

We return here to consistency of statements of the form $\chi \rightarrow [\theta]_{\sigma,2}^2$ (i.e. for every $c : [\chi]^2 \rightarrow \sigma$ there is $A \in [\chi]^\theta$ such that on $[A]^{\leq 2}$, c has at most two values), (when $2^{<\mu} \geq \chi > \theta^{>\mu}$, of course). In [Sh276, §2] this was done for $\mu = \aleph_0$, $\chi = 2^\mu$, $\theta = \aleph_1$ and χ quite large (in the original universe it is an Erdos cardinal). Originally, it was written for any $\mu = \mu^{<\mu}$ (χ measurable in the original universe) but because of the referee urging it is written up there for $\mu = \aleph_0$ only; though with an eye on the more general result which is only stated. In [Sh288] the main objective is to replace colouring of pairs by colouring of n -tuples (and even $(< \omega)$ -bytes). By [Sh284] we can lower the consistency strength. The main point of this section is to increase θ , and this time do it for $\mu = \mu^{>\mu} > \mu$, too.

2.1 Definition. A forcing notion Q satisfies $*_\mu^\epsilon$ where ϵ is a limit ordinal $< \mu$, if player I has a winning strategy in the following game $*_\mu^\epsilon[Q]$ defined as follows:
Playing: the play finishes after ϵ moves.

In the α -th the move:

Player I — if $\alpha \neq 0$ he chooses $\langle q_\zeta^\alpha : \zeta < \mu^+ \rangle$ such that $q_\zeta^\alpha \in Q$

and $(\forall \beta < \alpha)(\forall \zeta < \mu^+) p_\zeta^\beta \leq q_\zeta^\alpha$ and he chooses a regressive function $f_\alpha : \mu^+ \rightarrow \mu^+$ (i.e. $f_\alpha(i) < 1 + i$); if $\alpha = 0$ let $q_\zeta^\alpha = \emptyset_Q$, $f_\alpha = \emptyset$.

Player II — he chooses $\langle p_\zeta^\alpha : \zeta < \mu^+ \rangle$ such that $q_\zeta^\alpha \leq p_\zeta^\alpha \in Q$.

The Outcome: Player I wins provided whenever $\mu < \zeta < \xi < \mu^+$, $cf(\zeta) = cf(\xi) = \mu$ and $\bigwedge_{\beta < \epsilon} f_\beta(\zeta) = f_\beta(\xi)$ the set $\{p_\zeta^\alpha : \alpha < \epsilon\} \cup \{p_\xi^\alpha : \alpha < \epsilon\}$ has an upper bound in Q .

Remark. In the case $\mu = \aleph_0$ we can use Knaster condition instead.

2.2 Definition. Assume P, R are forcing notions, $P \subseteq R$, $P \triangleleft R$. Moreover, for simplicity, for $r \in R$ a member $r \upharpoonright P \in P$ is defined such that:

- (a) $r \upharpoonright P \leq r$,
- (b) $(\forall p)(r \upharpoonright P \leq p \in P \Rightarrow r)$, p compatible in R
- (c) $r^1 \leq r^2 \Rightarrow r^1 \upharpoonright P \leq r^2 \upharpoonright P$

(so $P \triangleleft R$).

Let (R, P) satisfies $*_\mu^\epsilon$ mean:

- (α) P satisfies $*_\mu^\epsilon$
- (β) If St_1 is a winning strategy for player I in the game $*_\mu^\epsilon[P]$ then in the following game called $*_\mu^\epsilon[P, R; St_1]$ the first player has a winning strategy:

Playing: as before but $\langle \langle q_\zeta^\theta \upharpoonright P : \zeta < \mu^+ \rangle, \langle p_\zeta^\theta \upharpoonright P : \zeta < \mu^+ \rangle : \alpha < \epsilon \rangle$ is required to be a play of $*_\mu^\epsilon[P]$ in which first player uses the strategy St_1 .

The outcome: Player I wins provided that: if $\zeta < \xi < \mu^+$, $cf(\zeta) = cf(\xi) = \mu$, $\bigwedge_{\beta < \alpha} f_\beta(\zeta) = f_\beta(\xi)$ and $r^* \in P$ is a $\leq_P$ -upper bound of $\{p_\zeta^\alpha \upharpoonright P : \alpha < \epsilon\} \cup \{p_\xi^\alpha \upharpoonright P : \alpha < \epsilon\}$, then $\{r\} \cup \{p_\zeta^\alpha : \alpha < \epsilon\} \cup \{p_\xi^\alpha : \alpha < \epsilon\}$ has an upper bound in R .

2.3 Definition/Lemma. Let $\mu = \mu^{<\kappa} < \kappa\sigma = cf\kappa \leq \lambda \leq \chi$. (Usually fixed hence suppressed in the notion). We define and prove the following by induction on (the ordinal) α :

1) [Def] We let $\mathcal{K}^{\epsilon,\alpha} = \mathcal{K}_{\mu,\kappa,\lambda,\chi}^{\epsilon,\alpha}$ is the family of sequences $\bar{Q} = \langle P_\beta, Q_\beta, a_\beta, \mathbf{t}_\beta : \beta < \alpha \rangle$ such that:

- (a) $\langle P_\beta, Q_\beta : \beta < \alpha \rangle$ is a $(< \mu)$ -support iteration (so $P_\alpha = \lim \bar{Q}$ denotes the natural limit)
- (b) $a_\beta \subseteq \beta$, $|a_\beta| < \kappa$, $[\gamma \in a_\beta \Rightarrow a_\gamma \subseteq a_\beta]$
- (c) Q_β has cardinality $< \lambda$ and is a P_{a_β} -name (see part (*) below)
- (d) $\mathbf{t}_\beta$ is a truth value; i.e. true or false
- (e) if $\mathbf{t}_\beta$ is true, $\beta < \gamma \leq \alpha$ then (P_γ, P_β) satisfies $*_\mu^\epsilon$.

2) [Def] For $\bar{Q}$ as above:

- (a) $a \subseteq \alpha$ is called $\bar{Q}$ -closed if $\beta \in a \Rightarrow a_\beta \subseteq a$
- (b) for a $\bar{Q}$ -closed subset a of α we let
$$P_a = \left\{ p \in P_\alpha : \text{Dom}(p) \subseteq a \text{ and for each } \beta \in \text{Dom}(p) \text{ we have } p(\beta) \right. \\ \left. \text{is a } P_{a \cap \beta}\text{-name} \right\}$$

$$P_a^* = \{ p \in P_\alpha : \text{Dom } p \subseteq a \text{ and for each } \beta \in \text{Dom}(p) \text{ we have } p(\beta) \text{ is a } P_{a_\beta}\text{-name} \\ \text{and if } Q_\beta \subseteq V \text{ then } p(\beta) \text{ is from } V \text{ (see 0.B.)} \}$$

3) [Lemma] For $\bar{Q}$ as above, $\beta < \alpha$

- (a) $\bar{Q} \upharpoonright \beta \in \mathcal{K}^{\epsilon,\beta}$
- (b) if $a \subseteq \beta$ then a is Q -closed iff a is $(\bar{Q} \upharpoonright \beta)$ -closed
- (c) if $a \subseteq \alpha$ is $\bar{Q}$ -closed then so is $a \cap \beta$.

4) [Lemma] For $\bar{Q}$ as above, and $p < \alpha$, $P_\beta < P_\alpha$, moreover, if $p \in P_\alpha$, $p \upharpoonright \beta \leq q \in P_\beta$ then $(p \upharpoonright [\beta \cup \alpha]) \cup q \in P_\alpha$ is a lub of p, q .

5) [Lemma] For $\bar{Q}$ as above

- (a) P_α^* is a dense subset of P_α
- (b) if a is $\bar{Q}$ -closed then $P_a < P_\alpha$, P_a^* is a dense subset of P_a .

6) If $a \subseteq b$ is a $\bar{Q}$ -closed set then (P_b^*, P_a^*) satisfies $*_\mu^\epsilon$.

7) The sequence $\bar{Q} = \langle P_\beta, Q_\beta, a_\beta, \mathbf{t}_\beta : \beta < \alpha \rangle$ belongs to $\mathcal{K}^{\epsilon,\mu}$ if α is a limit ordinal,

$\bigwedge_{\gamma < \alpha} \bar{Q} \upharpoonright \gamma \in \mathcal{K}^{\epsilon,\gamma}$ for some club C of α , $\bigwedge_{\gamma \in C} \mathbf{t}_\gamma = \text{true}$.

8) The sequence $\bar{Q} = \langle P_\beta, Q_\beta, a_\beta, \mathbf{t}_\beta : \beta < \alpha \rangle$ belongs to $\mathcal{K}^{\epsilon,\mu}$ if $\alpha = \gamma + 1$, $a \subseteq \gamma$ is a $(\bar{Q} \upharpoonright \gamma)$ -closed set of cardinality $< \kappa$, Q_β is a $P_{a_\beta}^*$ -name of a forcing notion satisfying $*_\mu^\epsilon$ and $\mathbf{t}_\beta = \text{true}$.

Proof. [Saharon].

2.4 Theorem. *Suppose $\mu = \mu^{<\mu} < \kappa = \lambda < \chi$, χ is measurable. For some forcing notion P of cardinality χ , μ -complete not collapsing cardinalities not changing cofinalities we have:*

$\Vdash_P "2^\mu = \chi \text{ and for every } \sigma < \chi \text{ and } \theta < \kappa \text{ we have } \chi \rightarrow [\theta]_{\sigma,2}^2"$ (and Axiom as in [Sh280]) (and the parallel of M for forcing notions satisfying $*_\mu^\epsilon$).

Proof. Fix $\epsilon < \mu$, $\mathcal{K}^\alpha = \mathcal{K}^{\epsilon,\alpha}$, $\mathcal{K} = \bigcup_{\alpha < \chi} \mathcal{K}^\alpha$. By preliminary forcing without loss of generality " χ measurable" is preserved by forcing with $(\chi^{>2}, \trianglelefteq)$, (= adding a Cohen subset of χ). Let us define a forcing notion R :

$R = \{\bar{Q} : \bar{Q} \in \mathcal{K} \text{ for some } \alpha < \chi \text{ and } \bar{Q} \in H(\chi)\}$
ordered by: $\bar{Q}^1 \subseteq \bar{Q}^2$ iff $\bar{Q}^1 = \bar{Q}^2 \restriction \ell g(\bar{Q}^1)$.

As R is equivalent to $(\chi^{>2}, \trianglelefteq)$ we know that in V^R , χ is still measurable. Let $\bar{Q}^\oplus = \langle P_\beta, Q_\beta, a_\beta, \mathbf{t}_\beta : \beta < \chi \rangle$ be $\cup G_R$ and P_χ be the limit so $P_\chi^* \subseteq P_\chi$ is a dense subset. Now $R * P_\chi^*$ is the forcing P we have promised. The non-obvious point is $\Vdash_{R * P_\chi^*} "\chi * [\theta]_{\sigma,2}^2$ (where $\theta < \chi, \sigma < \chi$). So suppose $(r^*, p) \in R * P_\chi^*$, $(r^*, p) \Vdash$ "the colouring $\tau : [\chi]^2 \rightarrow \sigma$ is a counterexample". Let $\chi_1 = (2^\chi)^+$. Let $G_R \subseteq R$ be generic over V , $r^* \in G_R$. By [Sh289],xx in V^R we can find $B \in [\chi]^\chi$ and $\langle M_s : s \in [B]^{<\aleph_0} \rangle$ and $\langle f_{s,t} : (s,t) \in \bigcup_{n < \omega} [B]^n \times [B]^n \rangle$ such that $M_s \prec (H(\chi_1)^{V[G_R]}, H(\chi_1), \in)$, $\{\chi, G_R, p, \tau\} \in M_s$, $M_s \cap M_t = M_{s \cap t}$, $[M_s]^{\leq (\kappa + \lambda + 2^\mu)} \subseteq M_s$, $f_{s,t}$ isomorphism from M_s onto M_t as there.

Let $C = \left\{ \delta < \chi : \delta = \sup(\beta \cap \delta) \text{ and } (s \in [\delta]^{<\aleph_0} \Rightarrow M_s \cap \chi \subseteq \delta) \right\}$.

Let $\gamma(*) = \text{Min}(B)$. Now for $p \in P^\oplus \cap M_{\gamma(*)}$ and $u = \{c_1, c_2\} \in [\sigma]^{\leq 2}$ let us define the statement

$(*)_p^u$: if $p \leq p^0 \in p^* \cap M_{\gamma(*)}$ then we can find $p^1, p^2 \in P^* \cap M_{\gamma(*)}$, $p^0 \leq p^1$, $p^0 \leq p^2$ such that:

for $\gamma_1 \neq \gamma_2$, $\gamma_2 \in B$, $\gamma_2 \in B$, we can find $r_1, r_2 \in P^* \cap M_{\{\gamma_1, \gamma_2\}}$ (so $\text{Dom } r_\ell \subseteq M_{\{\gamma_1, \gamma_2\}} \cap \chi$) such that:

$$r_\ell \Vdash "\tau(\{\gamma_1, \gamma_2\}) = c_\ell"$$

$$r_\ell \restriction (\lambda \cap M_{\{\gamma_\ell\}}) = f_{\{\gamma(*)\}, \{\gamma_\ell\}}(P^1)$$

$$r_\ell \restriction (\lambda \cap M_{\{\gamma_{3-\ell}\}}) = f_{\{\gamma(*)\}, \{\gamma_{3-\ell}\}}(P^2).$$

Easily $I = \{p \in P^* \cap M_{\{\gamma(*)\}} : \text{for some } u, (*)_p^u \text{ hold}\}$ is a dense subset of $P^* \cap M_{\{\gamma(*)\}}$, but this partial forcing satisfies the μ^+ -c.c. hence we can find $\mathbf{I}^* = \{p_\zeta : \zeta < \mu\} \subseteq I$ a maximal antichain of $P^* \cap M_{\{\gamma(*)\}}$ hence of P^* , and let

$$(*)_\beta^{(c_1[p], c_2[p])} \quad \text{for } p \in \mathbf{I}^*.$$

As G_R was any generic subset of R to which r^* belongs we have R -names $\gamma(*) \langle \langle p_\zeta, c_1(p_\zeta), c_2(p_\zeta) \rangle : \zeta < \mu \rangle, \langle \underline{M}_s : s \in [\underline{B}]^{<\aleph_0} \rangle, \langle \underline{f}_{s,t} : (s,t) \in \bigcup_{n < \omega} [\underline{B}]^n \times [\underline{B}]^n \rangle$ forced by r^* to be as above. Without loss of generality r^* force a values $\gamma(*) \langle \langle p_\zeta^*, c_1^*(p_\zeta), c_2^*(p_\zeta) \rangle : \zeta < \mu \rangle, M_\emptyset, M_{\{\gamma(*)\}}, \langle \langle p_\zeta^*, c_1^*(p_\zeta), c_2^*(p_\zeta) \rangle : \zeta < \mu \rangle$.

We now choose by induction on $\zeta \leq \sigma + 1$, $Q^\zeta, \alpha^\zeta, \gamma^\zeta$ such that:

- (A)(a) $Q^\zeta \in R$
- (b) $\bar{Q}^0 = r^*$
- (c) $\ell g(\bar{Q}^\zeta) = \alpha^\zeta$
- (d) $\xi < \zeta \Rightarrow \bar{Q}^\xi = \bar{Q}^\zeta \restriction \alpha^\xi$
- (1) (e)" $\langle \alpha^\zeta : \zeta \leq \sigma + 1 \rangle$ is strictly increasing continuous
- (f) $\alpha^\zeta < \gamma^\zeta < \alpha^{\zeta+1}$
- (g) $\bar{Q}^{\zeta+1} \Vdash_R \text{"}\gamma^\zeta \in \underline{B}\text{"}$
- (h) $\bar{Q}^{\zeta+1}$ forces a value to $\langle \underline{M}_s : s \in [\underline{B} \cap (\gamma^\zeta + 1)^{<\omega}] \rangle = \langle \underline{M}_s : s \in [\underline{B}_\zeta]^{<\omega} \rangle$
- (B)(a) if $\zeta \leq \sigma + 1, cf(\zeta) > \mu$ then:
 - (a) $a_\zeta^{\bar{Q}^{\zeta+1}} = \cup \{ \chi \cap M_{\{\xi_1, \xi_2\}} : \{\xi_1, \xi_2\} \in [\{\gamma_\epsilon : \epsilon < \zeta\}]^{\leq 2} \}$
 - (b) $Q_{\alpha^\zeta}^{\bar{Q}^{\zeta+1}} = \{h : h \text{ a function } \text{Dom}(h) \in [\{\gamma_\epsilon : \epsilon < \zeta\}]^{<\mu}, \text{Rang } h \subseteq \mu\}$.

For $\gamma_\epsilon \in \text{Dom } h$, $f_{\{\gamma(*)\}, \{\gamma_\epsilon\}}(p_{h(\gamma_\epsilon)}^*) \in G_{P_\zeta}$ for $\epsilon_1 < \epsilon_2 < \zeta$, we can find p^1, p^2, r_1, r_2 such that: $p_\zeta^* \leq p^\ell \in M_{\{\eta(*)\}} \cap P_\chi^*$,
 $r_\ell \in P_\chi^* \cap M_{\{\gamma_{\epsilon_1}, \gamma_{\epsilon_2}\}}$,
 $r_\ell \restriction (\lambda \cap M_{\{\gamma_{\epsilon_\ell}\}}) = f_{\{\gamma(*)\}, \{\gamma_{\epsilon_1}\}}(p^1)$,
 $r_\ell \restriction (\lambda \cap M_{\{\gamma_{\epsilon_{3-\ell}}\}}) = f_{\{\gamma(*)\}, \{\gamma_{\epsilon_{3-\ell}}\}}(p^2)$,
 $r_1 \in G_{P_\zeta}$ or $r_2 \in G_{P_\zeta}$.

The point is to verify that defining $Q_{\alpha^\zeta}^{\bar{Q}^{\zeta+1}}$ as we do in clause (B)(a) is allowable; i.e. that the condition concerning $*_\mu^\epsilon$ from Definition 2.2 holds. Saharon more.

2.5 Remark. We can in 2.4 replace “measurable”, by k_1^2 -Mahlo, but it is not straightforward; .e.g. we may use the following version of squared demand.

Let χ be k -Mahlo, $\kappa < \chi$

$(*)_{\kappa}^{\kappa, k}$ there is $\bar{A} = \langle A_\alpha : \alpha < \chi \rangle$ and $\bar{C} = \langle C_\alpha : \alpha < S \rangle$ such that:

- (a) $S \subseteq \{\delta < \chi : cf\delta \leq \kappa\}$, $\{\delta \in S : cf\delta = \kappa\}$ is a stationary subset of χ
- (b) $C_\alpha \subseteq \alpha$, $[\beta \in c_\alpha \Rightarrow c_\beta = c_\alpha \cap \beta]$, otp $c_\alpha \leq \kappa$, c_α a club of α
- (c) $A_\alpha \subseteq \alpha$,
- (d) $\beta \in c_\alpha \Rightarrow A_\beta = A_\alpha \cap \beta$
- (e) $\{\lambda < \chi : \lambda \text{ inaccessible, and for every } x \subseteq \lambda \{\alpha < \chi : cf\alpha = \kappa < \alpha, x \cap \alpha = A_\alpha\} \text{ is stationary}\}$ not only is stationary but is k -Mahlo.

This can be obtained e.g. by iteration with Easton support, in which for each inaccessible we add $\bar{A}$, $\bar{c}$ satisfying (a)-(d) above, each condition being an initial segment.

* * *

We do now what we can for colouring of triples and more.
By [Sh 288],§4 we can deal with μ supercompact and, of course, with $\mu = \aleph_0$.

§3

3.1 Content. τ is a vocabulary = set of predicates including $<$, K^τ the class of τ -structures, μ such that $<^\mu$ is a (linear) well ordering, $M, N \in K^\tau$ are compatible if $M \upharpoonright (|M| \cap |N|) = N \upharpoonright (|M| \cap |N|)$, and the $M \cup N$ is $(|M| \cup |N|, \dots, R^M \cup R^N, \dots)_{R \in \tau}$. Universe of a model M is $|M|$ so for $M \in \tau$, $M = (|M|, \dots, R^M, \dots)_{R \in \tau}$, $M \in K$ is standard is $|M|$ is a set of order, $<^M$ usual ordering, $\tau = \tau \setminus \{<\}$, $M^- = M \upharpoonright \tau^-$.

3.2 Definition. 1) We call $M \in K^\tau$ 2-connected if for no $N_1 N_2$ do we have $N_1 \subseteq M$, $N_2 \subseteq M$, $M^- = N_1^- \cup N_2^-$ and $\|N_1 \cap N_2\| \leq 1$.
 2) $\text{cont}^\kappa(M) = \{A \subseteq |M| : A \subseteq M, \text{otp}(A) < \kappa, M \upharpoonright A \text{ is 2-connected}\}$,
 $\text{cont}^\kappa \cong (M) = \{(M \upharpoonright A) / \cong : A \subseteq |M|, |A| < \kappa, M \upharpoonright A \text{ is 2-connected}\}$.
 3) For D a set of members of K^τ , $D / \cong$ is $\{M / \cong : M \in D\}$.
 4) If D is a set of 2-connected members of K^τ , $\sigma < \kappa < \infty$, $K_\kappa^\tau[D] = \{M \in K^\tau : \text{cont}_\kappa \cong (M) \subseteq D / \cong\}$
 (so without loss of generality $(*)[M \in N \Rightarrow \|M\| < \kappa][A \subseteq M \in D \ \& \ M \upharpoonright A \text{ 2-connected} \Rightarrow M \upharpoonright A \in D]$.
 5) $M \in K_\tau$ is m -indecomposable if for every $x, y \in M$ there is A , $\{x, y\} \subseteq A \subseteq M$, $|A| < m$, $M \upharpoonright A$ 2-connected.
 6) $\text{id com}_{m,\kappa}(M) = \{A \subseteq M : |A| < \kappa \text{ and } M \upharpoonright A \text{ is } m\text{-indecomposable}\}$
 $\text{id com}_{m,\kappa} \cong |M| = \{(M \upharpoonright A) / \cong : A \in \text{id com}_{m,\kappa}(M)\}$.
 If $\kappa > \|M\|$ we omit it.

3.3 Theorem. Suppose

- (a) $\mu < \lambda, \mu = \mu^{<\mu}, \lambda$ measurable, θ
- (b) $1 < m < \omega, D$ a set of 2-connected members of K^τ satisfying $(*)_m$ of 2-2,
 $|\tau|$
- (c) $M \in K_k^\tau[D], \|M\| < \mu$ and $|M|$ is $\alpha^* \leq \mu$.

Then for some P and $\dot{N}$:

- (A) P is a μ -complete λ -c.c. forcing notion of cardinality λ
- (B) $\dot{N}$ is a P -name of a standard member of $K_k^\tau[D]$ with universe λ
- (C) $\dot{N}$ “if $c : {}^\omega \lambda * \mu$ then for some $A \subseteq \lambda$:
 $\text{otp } A = \alpha^*, \dot{N} \upharpoonright A \cong \mu$ and $(*)$ if $n < \omega$,
 $\alpha_0 < \alpha_1 < \dots < \alpha_{n-1} < \alpha_n$ are from A ,
 $\dot{N} \upharpoonright \{\alpha_0, \alpha_1, \dots, \alpha_{n-2}, \alpha_{n-1}\} \cong \dot{N} \upharpoonright \{\alpha_0, \alpha_1, \dots, \alpha_{n-2}, \alpha_n\}$ then
 $c(\{\alpha_0, \dots, \alpha_{n-2}, \alpha_{n-1}\}) = c(\{\alpha_0, \dots, \alpha_{n-2}, \alpha_n\})$ ”
- (D) $\Vdash_P$ “ $\text{id cont}_m \cong (\dot{N}) \subseteq \text{id cont}_m \cong (G)$ ”
- (E) $\Vdash_P$ “ $\lambda = \mu^+$ ”.

Proof. :

A Stage: The forcing:

- (α) Let Q be the set of $p = (s^p, N^p, \varphi^p) = (s, N, \varphi)$ such that
- (a) $s \subseteq \lambda$, $(s) < \mu$
 - (b) $N^p \in K_m^\tau[D]$ has universe ${}^k s = \{\langle \alpha_0, \dots, \alpha_{m-1} \rangle : \alpha_0 > \alpha_1 > \dots > \alpha_{m-1}\}$
 - (c) $<^{N^p}$ is the lexicographic order on ${}^k s$
 - (d) for $R \in \tau^-$ if $\langle \bar{\alpha}^0, \dots, \bar{\alpha}^{m-1} \rangle \in R^{N^p}$ then for $\ell \neq k (< n)$ $\bar{\alpha}^\ell \sim \bar{\alpha}^m$ which means $\alpha_0^\ell > \alpha_0^m > \alpha_1^\ell > \alpha_1^m > \dots$ or $\alpha_0^m > \alpha_0^\ell > \alpha_1^m > \alpha_1^\ell > \dots$. We shall not strictly distinguish between N^p and $N^p \upharpoonright \tau^-$
 - (e) φ is a function, $\text{Dom } \varphi = \text{id cont}_{m,\kappa}(M)$ and $\varphi(N \upharpoonright A)$ is an embedding of $N \upharpoonright A$ into M
 - (f) if $A, B \subseteq \text{Dom } \varphi$, $A \subseteq B$ moreover A is an initial segment of B by $<^N$ then $\varphi(N \upharpoonright A) \subseteq \varphi(N \upharpoonright B)$
 - (g) for $\alpha < \lambda$, $p \upharpoonright \alpha \leq q$ where $p \upharpoonright \alpha = (s^p \cap \alpha, N^p \upharpoonright {}^m(s^p \cap \alpha), \varphi^p \upharpoonright \text{id cont}_m(N^p \upharpoonright {}^m(s^p \cap \alpha)))$, and $\leq$ is defined in (β) below.
- (β) The order on $Q : p \leq q$ iff:
- (a) $s^p \subseteq s^q$
 - (b) $N^p \subseteq N^q$
 - (c) $\varphi^p \subseteq \varphi^q$
 - (d) if $A \subseteq \text{cont}_m(N^q)$, $|A \cap N^p| > 1$ then $A \subseteq N^p$
 - (e) if $A \subseteq N^q$, $|A| < m$, $|A \cap N^p| > 1$, $A \subseteq N^p$ then we can find A_1, A_2 such that $A = A_1 \cup A_2$, $|A_1 \cap A_2| \leq 1$, $A_1 \cap A_2 \subseteq N^p$, $N^p \upharpoonright A \upharpoonright \tau^- = N^p \upharpoonright A_1 \upharpoonright \tau^- \cup N^p \upharpoonright A_2 \upharpoonright \tau^-$
 - (f) if $B \in \text{Dom } \varphi^q$, $A = B \cap N^p \in \text{Dom } \varphi^p$ then B is an end extension of A .

Lastly, $P = Q \times \text{Levy}(\mu_2 < \lambda)$. But to simplify presentation we prove only that Q satisfies (A)-(D) (waiving (E)).

B Stage: Basic properties of the forcing notion:

- (α) if $p \leq q$ in Q , $A \subseteq N^q$, $|A| < m$, then we can find $A_\ell (\ell < \kappa)$ such that:
 “ $\langle A_\ell \setminus N^p : \ell < \kappa \rangle$ is a partition of $A \setminus N^p$
 $|A_\ell \cap N^p| \leq 1$
 $N^q \upharpoonright A = N^q \upharpoonright A_0 \upharpoonright \tau^- + N^q \upharpoonright A_1 \upharpoonright \tau^- + \dots + N^q \upharpoonright A_{k-1} \upharpoonright \tau^- + N^q \upharpoonright (A \cap N^p)$
 [why? by induction on $|A|$ using (A)(B)(e)].
- (β) if $p \leq q$ in Q , $B \in \text{id cont}_m(q)$ then $B \cap N^p \in \text{id com}_m(q)$
 [why? if $x, y \in B \cap N^p$ there is $A \subseteq B$, $|A| < m$, $\{x, y\} \subseteq A$, $M \upharpoonright A$ 2-connected, by (α) above necessarily $A \subseteq N^p$, so we are done].
- (γ) $(\theta, \leq)$ is transitive.
 [why? check $A(\beta)(a)$, (b), (c), (d). As for (A)(β)(e) use (α) above and as for (A)((β)(f) use (β) above].
- (δ) Assume $q \in Q$, $\alpha < \beta < \lambda$. Then $(p \upharpoonright \beta) \upharpoonright \alpha = p \upharpoonright \alpha \leq p \upharpoonright \beta \leq q$.
 [why? by (A)(α)(g)].
- (ϵ) if $p \leq q$ and $\alpha < \lambda$ then $p \upharpoonright \alpha \leq q \upharpoonright \alpha$.
 [why? check].
- (ζ) Assume $p_\ell \in Q (\ell = 0, 1, 2)$ $p_0 \leq p_1$, $p_0 \leq p_2$, $s^{p_0} = s^{p_1} \cap s^{p_2}$ and $q = (s^q, N^q, \varphi^q)$, $s^q = s^p \cup s^{p_2}$, $N^q \upharpoonright \tau^- = N^{p_1} \upharpoonright \tau^- \cup N^{p_2} \upharpoonright \tau^-$ and $\varphi^{p_1} =$

$$\varphi^{p_1} \cup \varphi^{p_2}.$$

Then $p_\ell \leq q \in Q$.

[why? we prove it by induction on $\sup(s^q)$].

First check $q \in Q$; i.e. $(A)(\alpha)$, now $(A)(\alpha)(a)$, (b) , (c) , (d) should be clear. For $(A)(\alpha)(d)$ we need:

$$(*)_0 \text{ id cont}_m(N^q) = \text{id cont}_m(N^{p_1}) \cup \text{id cont}_m(N^{p_2}).$$

For this it suffices to:

$(*)_2$ if $A \in \text{id cont}_m(N^r)$ then $A \subseteq N^{p_1}$ or $A \subseteq N^{p_2}$ which holds as if $x \in A \setminus N^{p_1}$, $y \in A \setminus N^{p_2}$, there is B , $\{x, y\} \in B \subseteq A$, $|B| < m$, $N^r \restriction B$ is 2-connected, but by (α) above applied twice we get contradiction.

We have shown

$$(*)_3 \text{ if } A \in \text{cont}_m(N^r), \text{ then } A \subseteq N^{p_1} \text{ or } A \subseteq N^{p_2}.$$

Now $(A)(\alpha)(f)$ is easy, as for $(A)(\alpha)(g) : (A)(\beta)(a)$, (b) , (c) are easy. $(A)(B)(d)$ holds by $(*)_3$ above. As for $(A)(\beta)(e)$, let $B \subseteq N^q$, $|B| < m$. By (α) above applied to $p_1 \restriction \alpha \leq p_1$ we can find $\langle x_\ell, A_\ell : \ell < \ell^* \rangle$, $N^{p_1} \restriction A_\ell$ 2-connected, $A_\ell \cap N^{p_1 \restriction \alpha} = \{x_\ell\}$, $N^{p_1} \restriction A = \sum_\ell N^{p_1} \restriction A_\ell + N^{p_1} \restriction (A \cap N^{p_1 \restriction \alpha})$.

For each ℓ if $A_\ell \cap N^{p_0} \dots$

Those notes were intended to give consistency results of the form. We have $k < \omega$ a class K of 1 model with set of finite submodels pregiven (closed under edgeless amalgamation) and set of forbidden large k -connected in appropriate sense submodels, we can have $\forall \kappa \forall n \forall M \in K, M \rightarrow (N)_\kappa^n$.

But the conditions on the set of finite submodels do not match the forcing construction such that appropriate amalgamations hold. For “no odd circle of small length” fine should be rephrased.

PRIVATE APPENDIX

(17.4.94): 1) Replace $\aleph_0$ by higher...? 2) Replace Ramsey by weakly compact??

(4.5.94:2) Remember [Sh 481], §2 make sense: we use the parallel of squared scales - have start lecture and and finish correction §1, correcting §2. Enough for $2^\mu \rightarrow [\mu^{++}]_3^2$.

Write down:

- (a) Mahlo (use Easton support instead R , (which use $(< \chi)$ -support, for a larger cardinal) (generic from system $\langle N_s : i \rangle$?? preserve or only continuity (see (c)(b);
- (b) check colouring
- (c) Try:
 - (α) some $\chi' < \chi, \chi' \rightarrow [\mu^+]_3^2$, and $\chi' < \mu^{+\omega}$ (so here the guessing of $\langle M_{\gamma_\varepsilon} : \varepsilon \leq \theta \rangle$ is not by initial segments but by $\langle M_{\gamma_\varepsilon} \cap \delta_i : \varepsilon \leq \theta \rangle$ or
 - (β) back to what I thought last week, trying to force almost disjoint set [Baumgartner]

(d) 289, §5 in ZFC?

5.5.94 Will check how preliminary forcing affect the systems of [Sh 289], I though (April) will be like tree, so enough, have to check) the proof of 481, §1 indicates the Knaster in [Sh288] can be weakened sometimes.

11.5.94 In 1.7; the April idea of forcing almost disjoint sets may help to $2^{\aleph_0}$ -c.c. $\times \aleph_2$ -c.c. $\Rightarrow 2^{\aleph_0}$ -c.c.

The systems we need sometimes are just names for $\underline{c}(s)$: this is less: when it is not sufficient? Different numbers.

ASSIGNMENTS FROM '92

- 0) explain for $\mu = \aleph_0$ in §2
- 1) 2-3 fil
- 2) 2-4 fil
- 3) 2-5 fil
- 4) 2-6 fil
- 5) explicit c.c.c. condition — easier if we represent $\beta(i)$ as a set of ordinals
 - (5A) $2^{\aleph_0} < \aleph_\omega$
 - (5B) $\chi \rightarrow [\theta]_{\sigma,2}^2$ for every $\theta; \sigma < \chi$
- 6) $\chi < 2^\mu??$
- 7) The reference of $\lambda \rightarrow [\mu]_\mu^3$ if $2^{<\mu} < \lambda \leq 2^\mu$ or so
- 8) Concerning §1, if $2^{\aleph_0}$ is successor of regular?
- 9) Can you guess square domain of $\langle s_u : u \in [\mu^+]^{\leq 2} \rangle |s_4| \subseteq \mu$ with $\chi < \mu^{+\omega}??$